

designed to create the protective  
coating on the damaged skin  
and prevent the bacterial biofilms  
formation

# ENTOMIX®

contains a composition of FLIP7  
antimicrobial peptides

a new look at the antibiotic  
resistance problem

## Method of application of **ENTOMIX®**

Before the application of **ENTOMIX®** necrotic elements in the wound should be removed. For the best effect the **ENTOMIX®** hydrogel coating must be applied directly to the surface of the wound, since the gel components must be in direct contact with bacteria.

**ENTOMIX®** should be applied directly to the damaged skin surface with a thin layer (up to 1 mm) at least 1 time per day.

In case of burns and diabetic ulcers hydrogel need to be covered with secondary dressings that do not contain oxidizing agents, proteolytic enzymes or cytotoxic antiseptics about 3 minutes after application of **ENTOMIX®** to the wound.

**ENTOMIX®** hydrogel coating can be used individually or as an adjuvant to increase the effectiveness of antibiotic treatment.

Photographic materials provided by customers, Clinical Hospital № 122 named after L.G. Sokolov and I. I. Dzhanelidze Research Institute of Emergency Medicine in Saint-Petersburg.

## DIABETIC ULCER



Before using **ENTOMIX®**

The result of using **ENTOMIX®**

## DIABETIC ULCER



After 1 month of using **ENTOMIX®**

## DIABETIC ULCER



Before using **ENTOMIX®**

The result of using **ENTOMIX®**

## DIABETIC ULCER



After 1 week of using **ENTOMIX®**

## DIABETIC ULCER



Before using **ENTOMIX<sup>®</sup>**

The result of using **ENTOMIX®**

## DIABETIC ULCER



After 1 week of using **ENTOMIX®**



## MICROBIAL ECZEMA



Before using **ENTOMIX**<sup>®</sup>

The result of using **ENTOMIX®**

## MICROBIAL ECZEMA



After 1 week of using **ENTOMIX®**

## TROPHIC ULCER



Before using **ENTOMIX®**

The result of using **ENTOMIX®**

## TROPHIC ULCER



After 2 months of using **ENTOMIX®**

## BURNS



Before using **ENTOMIX®**

The result of using **ENTOMIX®**

## BURNS



After 1 week of using **ENTOMIX®**

## BURNS



Before using **ENTOMIX<sup>®</sup>**

The result of using **ENTOMIX®**

## BURNS



After 1 week of using **ENTOMIX®**



## PYODERMA



Before using **ENTOMIX®**

The result of using **ENTOMIX®**

**PYODERMA**



After 1 week of using **ENTOMIX®**

## ANGULAR CHEILITIS (STOMATITIS)



Before using **ENTOMIX®**

The result of using **ENTOMIX®**

**ANGULAR CHEILITIS  
(STOMATITIS)**



After 3 days of using **ENTOMIX®**



Biotechnological company "Allofarm"  
199178 St. Petersburg,  
7th line V.O., d.76, lit. PROPERTY

+7(812)320-55-42  
+7(931)592-33-75

[www.entomix.ru](http://www.entomix.ru)  
[www.allofarm.ru](http://www.allofarm.ru)  
[allofarmspb@gmail.com](mailto:allofarmspb@gmail.com)

**Thank you for your attention!**

Store in a dry place at temperatures between + 2 ° C and + 8 ° C. Expiration date 24 months. TP TC

009/2011 TU 9158-006-72500079-2011 (rev. 1-4)